



Stink bugs (families Acanthosomatidae and Pentatomidae) feed on plant sap (Acanthosomatidae, Pentatominae) or blood (hemolymph) of their insect preys (Asopinae) with their piercing-sucking mouthparts. Fifty-five species are known to occur in the state of Idaho, illustrated below with their body length ranges. They include a number of pests (●) of cultivated plants.

## ACANTHOSOMATIDAE



*Elasmostethus cruciatus* (8-11 mm)  
*Elasmucha lateralis* (6-9 mm)



*Apoecilus bracteatus* (13-17 mm)



*Perillus bioculatus* (8.5-11.5 mm)



*Perillus bioculatus* (8.5-11.5 mm)



*Perillus exaptus* (5-8 mm)



*Perillus exaptus* (5-8 mm)



*Perillus lunatus* (7-8 mm)



*Perillus splendidus* (8-9 mm)



*Podisus brevispinus* (8-12 mm)



*Podisus maculiventris* (9.5-14 mm)



*Podisus placidus* (8-12 mm)



*Podisus sericeiventris* (8.5-12 mm)



*Zicrona americana* (4.5-7.5 mm)

## PENTATOMIDAE: ASOPINAE

## PENTATOMIDAE: PENTATOMINAE



*Aelia americana* (7-9 mm)



*Antheminia eurynota remota* (9-10 mm)



*Banasa dimidiata* (7-10 mm)



*Banasa euchlora* (8-11 mm)



*Banasa sordida* (10-11.5 mm)



*Banasa tumidifrons* (7-9 mm)



*Brochymena affinis* (14-18 mm)



*Brochymena quadripustulata* (13-18 mm)



*Brochymena sulcata* (12-16 mm)



*Chinavia hilaris* (13-19 mm)



*Chlorochroa congrua* (9-11 mm)



*Chlorochroa faceta* (9-11 mm)



*Chlorochroa granulosa* (11-13 mm)



*Chlorochroa ligata* (13-19 mm)



*Chlorochroa opuntiae* (11-16 mm)



*Chlorochroa sayi* (10-16 mm)



*Chlorochroa uhleri* (12-16 mm)



*Coenus delius* (8.5-10.5 mm)



*Cosmopepla conspicularis* (6-8 mm)



*Cosmopepla intergressa* (4-6 mm)



*Cosmopepla lintneriana* (4-7 mm)



*Cosmopepla uhleri* (4-6 mm)



*Dendrocoris pini* (5.5-7 mm)



*Euschistus conspersus* (11-12 mm)



*Euschistus inflatus* (11-13 mm)



*Euschistus variolarius* (11-15 mm)



*Halyomorpha halys* (14-17 mm)



*Holcostethus abbreviatus* (8-9.5 mm)



*Holcostethus limbolaris* (7-9 mm)



*Holcostethus tristis* (7.5-8.5 mm)



*Neottiglossa trilineata* (5-6 mm)



*Neottiglossa tumidifrons* (4-5 mm)



*Neottiglossa undata* (4.5-6 mm)



*Prionosoma podopoides* (8.5-11 mm)



*Tepa brevis* (6 mm)



*Tepa rugulosa* (5-7 mm)



*Tepa vanduzeei* (5-8 mm)



*Tepa yerma* (5.5-7 mm)



*Thyanta pallidivirens* (9-11 mm)



*Thyanta pallidivirens* (9-11 mm)



*Trichopepla atricornis* (6-8 mm)



*Trichopepla aurora* (8.5-9 mm)



*Trichopepla dubia* (6.5-8 mm)



*Trichopepla grossa* (9-10 mm)

● = Pest species

The brown marmorated stink bug (*Halyomorpha halys*) is the most severely damaging pest species, feeding on over 175 host plants in the U.S.A. Native to Asia, it was detected in Pennsylvania in 1996 and has since gradually spread westward, reaching Oregon in 2004 and Idaho in 2012.

© Copyright 2025 University of Idaho Barr Entomological Museum. Downloading, printing, sharing and distribution are allowed. Partial reproduction for non-commercial purposes allowed if original source is acknowledged. Unauthorized sale is strictly prohibited. This poster was compiled by Devolet T. Mtui and Luc Leblanc. All photographed specimens are digitized (<https://ecdysis.org>) and deposited at the William F. Barr Entomological Museum (University of Idaho, Moscow, ID).