Vegetable & Flower Planting Timetable and Seed Starting Plan

(All seeded, except for transplants labeled TP)



1250 W Iron wood, Ste 107 Coeur d°Alene, ID 83814

Phone: (208) 292-2525 Plant Clinic: (208)292-1377

E-mail: kootenaimg@uidaho.edu Web: uidaho.edu/kootenai



Indoors

Outdoors

	Under Lights	In Cold Frame
Early February	Onions, leeks, geraniums	
Mid-March	Eggplants, peppers, tomatoes	Broccoli, cauliflower,
	basil, parsley	cabbage,
	Verbena, petunias, impatiens,	Lettuce, spinach,
	coleus, alyssum, lobelia	mesclun, mache
Early April	Asters, zinnias, marigolds	In Garden
		Lettuce, spinach, arugula
Mid-April		Onion set, onion plants,
		seed potatoes, peas
Late April	Pak choi	Carrots, beets, chard,
		broccoli, cauliflower,
		cabbage (TP)
Early May	Squashes, melons,	Leeks (TP)
	cucumbers, pumpkins	
Mid-May		Parsnips, parsley (TP)
		Beans, corn
Late May		Tomato (TP) pak choi
		(TP)
Early June	Brussels sprouts	Pepper, eggplant, basil,
		cucumber, melon,
		squash
		(all from TPs)
Mid-July		Brussels sprouts (TP)
		2 nd crops—broccoli,
		beets, carrots, lettuce,
		pak choi
Early August		Kale, turnips
		2 nd cropspinach
Early October		Garlic, shallots,
		"walking" onions
Late October		In Cold Frame
Early November		Spinach, mache

YOUR SEED-STARTING PLAN The Spring Frost-Free Date in My Garden is_____ WHEN TO OPTIMUM SOIL WEEKS SAFE TO SET OUT TIME TEMP. FOR **CROP START** FROM (RELATIVE TO FROST-SETTING OUT DATE **GERMINATION INSIDE** SOWING FREE DATE) (in F) 70° 1 week after Basil 55°- 60° Beets* 4-6 2 weeks before 60° Broccoli 4-6 2 weeks before 60° Cabbage 4-6 4 weeks before Cauliflower 60° 2 weeks before 4-6 45°-80° Collards 4-6 4 weeks before 70°-80° 3-4 1 to 2 weeks after Cucumber 80° **Eggplant** 8-10 2 to 3 weeks after 60° Kale 4-6 4 weeks before Kohlrabi* 80° (optimal) 4-6 4 weeks before 55°-75° 3 to 4 weeks before Lettuce 4-5 70°- 75° Melons 3-4 2 weeks after Mustard* 45°-70° 4-6 4 weeks before 55° Onions 6-8 4 weeks before 70° 9-10 2 to 3 weeks before Parsley 60° Peas* 3-4 6 to 8 weeks before 70° 6-14 2 weeks after <u>Peppers</u> Pumpkins 65° 3-4 2 weeks after 60° Spinach 4-6 3 to 6 weeks before 65° Squash 3-4 2 weeks after 55°-60° Swiss chard 4-6 2 weeks before 70° 6-8 <u>Tomatoes</u> 1 to 2 weeks after * These crops are usually direct seeded outdoors, but they can be started inside.